

LA-UR- 03 -8908

Approved for public release;
distribution is unlimited.

Title: Hydrodynamic Models of the Accretion Stream in
Magnetic Cataclysmic Variables

Author(s): Kunegunda E. Belle

Submitted to: Proceedings of IAU Colloquium 194: Compact Binaries in
the Galaxy and Beyond



Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the University of California for the U.S. Department of Energy under contract W-7405-ENG-36. By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

Form 836 (8/00)



HYDRODYNAMIC MODELS OF THE ACCRETION STREAM IN MAGNETIC CATAclysmic VARIABLES

Kunegunda E. Belle¹

Here I use the MHD code, FLASH, from the University of Chicago ASCI Center for Astrophysical Thermonuclear Flashes, to simulate initial two-dimensional simple models of accretion streams in magnetic CVs. These models explore the evolution of inflowing material for two white dwarf masses; $0.5M_{\odot}$ and $0.9M_{\odot}$. It is seen that a discontinuity forms at a height of 6.1×10^8 cm above the white dwarf 'surface' for $M_{WD} = 0.5M_{\odot}$ and at 1.6×10^9 cm for $M_{WD} = 0.9M_{\odot}$. These models will be developed further with the ultimate goal of completing a three-dimensional model of magnetic accretion onto a WD surface.

The Models: Models using different white dwarf masses were constructed. For each model, a 2D cartesian grid of width 4×10^7 cm and height 5×10^9 cm was used. An accretion stream of this width roughly corresponds to a fractional accretion area of $f = 0.001$ on the surface of a $0.9M_{\odot}$ white dwarf. The computational grid is set up so that one computational block is 2×10^7 cm on a side and three levels of adaptive mesh refinement (AMR) are allowed during the computation. Protons and electrons are used for the stream composition, the material is allowed to radiate, and a gamma-law EOS and plane parallel gravity are employed. The entire computational volume is given the initial stream parameters $T = 50,000$ K and $\rho = 1 \times 10^{-10}$ g cm⁻³. A downward velocity of -3×10^5 cm s⁻¹ is given to material above $y = 3 \times 10^9$ cm. Each model uses a 1 MG magnetic field.

Model 1. This simulation was performed for $M_{WD} = 0.5M_{\odot}$ ($\log g = 7.86$). Figure 1 displays the temperature profile at times $t = 0$ and $t = 69$ s. This is the time at which a discontinuity forms at $y = 6.1 \times 10^8$ cm, as can be seen in the plot. After time $t = 69$ s, a numerically seeded instability develops.

Model 2. This simulation was performed for $M_{WD} = 0.9M_{\odot}$ ($\log g = 8.49$). Figure 2 displays the temperature profile at times $t = 0$ and $t = 42.5$ s. This is the time at which a discontinuity forms at




Fig. 1. Temperature profile for Model 1 ($M_{WD} = 0.5M_{\odot}$).




Fig. 2. Temperature profile for Model 2 ($M_{WD} = 0.9M_{\odot}$).

$y = 1.6 \times 10^9$ cm, as can be seen in the plot. After time $t = 42$ s, a numerically seeded instability develops.

The software used in this work was in part developed by the DOE-supported ASCI/Alliance Center for Astrophysical Thermonuclear Flashes at the University of Chicago.

¹X-2 MS P225, Los Alamos National Laboratory, Los Alamos, NM 87545 (belle@lanl.gov).